

## HEPATITIS C (HCV)/HIV COINFECTION (Updated December 1, 2009)

Long-term studies of patients with chronic hepatitis C virus (HCV) infection show that approximately 33% of the patients progress to cirrhosis at a median time of less than 20 years [1-2]. This rate of progression increases with older age, alcoholism, male sex, and HIV infection [3-6]. A meta-analysis demonstrated that the rate of progression to cirrhosis for persons coinfecting with HCV/HIV was about three times higher than the rate for HCV mono-infected patients [5]. This accelerated rate is magnified in patients with low CD4 counts. Chronic HCV infection also complicates HIV treatment due to the increased frequency of antiretroviral (ARV)-associated hepatotoxicity [7-8]. Multiple studies have shown poor prognosis for HCV/HIV coinfection in the era of combination antiretroviral therapy (ART). It is unclear if HCV infection accelerates the rate of HIV progression [9] or if the accelerated rate primarily reflects the impact of injection drug use, which is strongly linked to HCV infection [10-11]. Although whether ART reduces the attributable morbidity/mortality from untreated HCV is unknown, the presence of chronic HCV infection influences the treatment of HIV with ARV as discussed below.

### **Assessment of HCV/HIV Coinfection Prior to Antiretroviral Therapy**

- Prior to initiation of ART, HIV-infected patients should be screened for HCV infection with sensitive immunoassays licensed for detection of antibody to HCV in blood. To confirm the presence of chronic infection, HCV-seropositive persons should be tested for HCV RNA using a qualitative or quantitative assay [12].
- Patients with HCV/HIV coinfection should be advised to avoid alcohol consumption, use appropriate precautions to prevent transmission of both viruses to others, and receive hepatitis A (HAV) and hepatitis B (HBV) vaccines if susceptible.
- All patients with HCV/HIV coinfection should be evaluated for HCV therapy. HCV treatment is recommended according to standard guidelines with strong preference for treating patients with higher CD4 counts. For patients with lower CD4 counts (<200 cells/mm<sup>3</sup>), it may be preferable to initiate ART and delay HCV therapy until CD4 counts increase as a result of HIV treatment [12-15].
- Concurrent treatment of both HIV and HCV is feasible but may be complicated by pill burden, drug toxicities, and drug interactions. Some notable considerations include:
  - Didanosine (ddI) should not be given with ribavirin because of the potential for drug-drug interactions leading to life-threatening ddI-associated mitochondrial toxicity including hepatomegaly/steatosis, pancreatitis, and lactic acidosis [16].
  - Zidovudine (ZDV) combined with ribavirin should be avoided when possible because the higher rates of anemia associated with the combination make ribavirin dose reduction necessary [17].
  - Abacavir (ABC) has been associated with decreased response to peginterferon plus ribavirin in some, but not all, retrospective studies; current evidence is insufficient to recommend avoiding this combination [18-20].
  - Growth factors (e.g., filgrastim and erythropoietin) may be required to manage interferon-associated neutropenia and ribavirin-associated anemia; ZDV may increase the need for adjuvant growth factors due to increased bone marrow suppression [17].

### **Antiretroviral Therapy in HCV/HIV Coinfection**

- Hepatotoxicity: Drug-induced liver injury (DILI) following ART is more common in HIV/HCV coinfection. The greatest risk of DILI may be observed in coinfecting persons with advanced liver disease (e.g., cirrhosis or end-stage liver disease) [21]. Eradication of HCV infection may decrease the likelihood of ARV-associated DILI [22].
  - Given the substantial heterogeneity in patient populations and drug regimens, comparison of DILI incidence rates for individual ARV agents across clinical trials is difficult. In such studies, the highest incidence rates of Grade 3 or 4 elevations in liver enzyme levels have been observed during therapy with ARV drugs that are no longer commonly used in clinical practice and that include stavudine (d4T) (with or without ddI), nevirapine (NVP), full-dose ritonavir (RTV) (600 mg twice daily), or tipranavir (TPV) (boosted by low-dose RTV) [23]. Also, due to the potential for concurrent fatty liver disease (steatosis), the use of d4T or ddI should be limited [24].
  - Patients should be monitored by following alanine aminotransferase (ALT) and aspartate aminotransferase (AST) levels at 1 month and then every 3 months after initiation of ART. Mild to moderate fluctuations in ALT and/or AST are typical in persons with chronic HCV infection. In the

absence of signs and/or symptoms of liver disease these fluctuations do not require interruption of ART. Significant ALT and/or AST elevation (>5 times the upper limit of the laboratory reference range) should prompt careful evaluation for signs and symptoms of liver insufficiency and for alternative causes of liver injury (e.g., acute HAV or HBV infection, hepatobiliary disease, or alcoholic hepatitis); short-term interruption of ART may be required [25].

- When to start ART: The rate of liver disease (fibrosis) progression is accelerated by HIV/HCV coinfection, particularly in persons with low CD4 counts ( $\leq 350$  cells/mm<sup>3</sup>). Data derived largely from retrospective cohort studies regarding the effect of ART on the natural history of HCV disease are inconsistent [6, 26-27]. However, ART may slow the progression of liver disease by preserving or restoring immune function and reducing HIV-related immune activation and inflammation [28-30]. Thus, for most coinfecting patients including those with cirrhosis, the potential benefits of ART outweigh concerns regarding DILI.
  - ART should be started in HCV/HIV-coinfecting persons in accordance with the Panel's recommendation for initiating ART in ART-naïve patients.
- What to start and what not to use: Initial combination regimens for the ARV-naïve patient with HCV/HIV are the same as for persons without HCV infection. HCV infection does not significantly alter the virologic or immunologic response to effective ART [31]. Special considerations for ART in persons with HCV/HIV coinfection include:
  - Patients receiving or considering therapy with ribavirin should avoid ddI, d4T, and ZDV.
  - ARV agents with the greatest risk of DILI should be used with caution.
  - Cirrhotic patients should be carefully assessed for signs of liver decompensation according to the Child-Turcotte-Pugh classification system because hepatically metabolized ARV drugs may require dose modification or avoidance in patients with Child-Pugh class B and C disease. (See [Appendix B, Table 7.](#))

## References

1. Alter MJ, Margolis HS, Krawczynski K, et al. The natural history of community-acquired hepatitis C in the United States. The Sentinel Counties Chronic non-A, non-B Hepatitis Study Team. *N Engl J Med*. 1992;327(27):1899-1905.
2. Thomas DL, Astemborski J, Rai RM, et al. The natural history of hepatitis C virus infection: host, viral, and environmental factors. *JAMA*. 2000;284(4):450-456.
3. Poynard T, Bedossa P, Opolon P. Natural history of liver fibrosis progression in patients with chronic hepatitis C. The OBSVIRC, METAVIR, CLINIVIR, and DOSVIRC groups. *Lancet*. 1997;349(9055):825-832.
4. Wiley TE, McCarthy M, Breidi L, et al. Impact of alcohol on the histological and clinical progression of hepatitis C infection. *Hepatology*. 1998;28(3):805-809.
5. Graham CS, Baden LR, Yu E, et al. Influence of human immunodeficiency virus infection on the course of hepatitis C virus infection: a meta-analysis. *Clin Infect Dis*. 2001;33(4):562-569.
6. Thein HH, Yi Q, Dore GJ, et al. Natural history of hepatitis C virus infection in HIV-infected individuals and the impact of HIV in the era of highly active antiretroviral therapy: a meta-analysis. *AIDS*. 2008;22(15):1979-1991.
7. Sulkowski MS, Thomas DL, Chaisson RE, et al. Hepatotoxicity associated with antiretroviral therapy in adults infected with human immunodeficiency virus and the role of hepatitis C or B virus infection. *JAMA*. 2000;283(1):74-80.
8. Sulkowski MS, Thomas DL, Mehta SH, et al. Hepatotoxicity associated with nevirapine or efavirenz-containing antiretroviral therapy: role of hepatitis C and B infections. *Hepatology*. 2002;35(1):182-189.
9. Greub G, Ledergerber B, Battegay M, et al. Clinical progression, survival, and immune recovery during antiretroviral therapy in patients with HIV-1 and hepatitis C virus coinfection: the Swiss HIV Cohort Study. *Lancet*. 2000;356(9244):1800-1805.
10. Vlahov D, Graham N, Hoover D, et al. Prognostic indicators for AIDS and infectious disease death in HIV-infected injection drug users: plasma viral load and CD4+ cell count. *JAMA*. 1998;279(1):35-40.
11. Celentano DD, Vlahov D, Cohn S, et al. Self-reported antiretroviral therapy in injection drug users. *JAMA*. 1998;280(6):544-546.
12. Ghany MG, Strader DB, Thomas DL, et al. Diagnosis, management, and treatment of hepatitis C: an update. *Hepatology*. 2009;49(4):1335-1374.
13. Soriano V, Puoti M, Sulkowski M, et al. Care of patients coinfecting with HIV and hepatitis C virus: 2007 updated recommendations from the HCV-HIV International Panel. *AIDS*. 2007;21(9):1073-1089.
14. Tien PC. Management and treatment of hepatitis C virus infection in HIV-infected adults: recommendations from the Veterans Affairs Hepatitis C Resource Center Program and National Hepatitis C Program Office. *Am J Gastroenterol*. 2005;100(10):2338-2354.
15. Avidan NU, Goldstein D, Rozenberg L, et al. Hepatitis C Viral Kinetics During Treatment With Peg IFN-alpha-2b in HIV/HCV Coinfecting Patients as a Function of Baseline CD4+ T-Cell Counts. *J Acquir Immune Defic Syndr*. 2009.

16. Fleischer R, Boxwell D, Sherman KE. Nucleoside analogues and mitochondrial toxicity. *Clin Infect Dis*. 2004;38(8):e79-80.
17. Alvarez D, Dieterich DT, Brau N, et al. Zidovudine use but not weight-based ribavirin dosing impacts anaemia during HCV treatment in HIV-infected persons. *J Viral Hepat*. 2006;13(10):683-689.
18. Vispo E, Barreiro P, Pineda JA, et al. Low response to pegylated interferon plus ribavirin in HIV-infected patients with chronic hepatitis C treated with abacavir. *Antivir Ther*. 2008;13(3):429-437.
19. Laufer N, Laguno M, Perez I, et al. Abacavir does not influence the rate of virological response in HIV-HCV-coinfected patients treated with pegylated interferon and weight-adjusted ribavirin. *Antivir Ther*. 2008;13(7):953-957.
20. Mira JA, Lopez-Cortes LF, Barreiro P, et al. Efficacy of pegylated interferon plus ribavirin treatment in HIV/hepatitis C virus co-infected patients receiving abacavir plus lamivudine or tenofovir plus either lamivudine or emtricitabine as nucleoside analogue backbone. *J Antimicrob Chemother*. 2008;62(6):1365-1373.
21. Aranzabal L, Casado JL, Moya J, et al. Influence of liver fibrosis on highly active antiretroviral therapy-associated hepatotoxicity in patients with HIV and hepatitis C virus coinfection. *Clin Infect Dis*. 2005;40(4):588-593.
22. Labarga P, Soriano V, Vispo ME, et al. Hepatotoxicity of antiretroviral drugs is reduced after successful treatment of chronic hepatitis C in HIV-infected patients. *J Infect Dis*. 2007;196(5):670-676.
23. Nunez M. Hepatotoxicity of antiretrovirals: incidence, mechanisms and management. *J Hepatol*. 2006;44(1 Suppl):S132-139.
24. McGovern BH, Ditelberg JS, Taylor LE, et al. Hepatic steatosis is associated with fibrosis, nucleoside analogue use, and hepatitis C virus genotype 3 infection in HIV-seropositive patients. *Clin Infect Dis*. 2006;43(3):365-372.
25. Sulkowski MS, Thomas DL. Hepatitis C in the HIV-infected patient. *Clin Liver Dis*. 2003;7(1):179-194.
26. Sulkowski MS, Mehta SH, Torbenson MS, et al. Rapid fibrosis progression among HIV/hepatitis C virus-co-infected adults. *AIDS*. 2007;21(16):2209-2216.
27. Brau N, Salvatore M, Rios-Bedoya CF, et al. Slower fibrosis progression in HIV/HCV-coinfected patients with successful HIV suppression using antiretroviral therapy. *J Hepatol*. 2006;44(1):47-55.
28. Macias J, Berenguer J, Japon MA, et al. Fast fibrosis progression between repeated liver biopsies in patients coinfecting with human immunodeficiency virus/hepatitis C virus. *Hepatology*. 2009;50(4):1056-1063.
29. Verma S, Goldin RD, Main J. Hepatic steatosis in patients with HIV-Hepatitis C Virus coinfection: is it associated with antiretroviral therapy and more advanced hepatic fibrosis? *BMC Res Notes*. 2008;1:46.
30. Ragni MV, Nalesnik MA, Schillo R, et al. Highly active antiretroviral therapy improves ESLD-free survival in HIV-HCV co-infection. *Haemophilia*. 2009;15(2):552-558.
31. Miller MF, Haley C, Koziel MJ, et al. Impact of hepatitis C virus on immune restoration in HIV-infected patients who start highly active antiretroviral therapy: a meta-analysis. *Clin Infect Dis*. 2005;41(5):713-720.